



Calcium signaling and β 2-adrenergic receptors regulate 1-nitropyrene induced CXCL8 responses in BEAS-2B cells.

Abdullah Mayati, E. Le Ferrec, Jørn A Holme, Olivier Fardel, Dominique Lagadic-Gossmann, Johan Ovrevik

► To cite this version:

Abdullah Mayati, E. Le Ferrec, Jørn A Holme, Olivier Fardel, Dominique Lagadic-Gossmann, et al.. Calcium signaling and β 2-adrenergic receptors regulate 1-nitropyrene induced CXCL8 responses in BEAS-2B cells.. *Toxicology in Vitro*, 2014, 28 (6), pp.1153-7. 10.1016/j.tiv.2014.05.012 . hal-01058607

HAL Id: hal-01058607

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01058607>

Submitted on 7 Oct 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Title:

2 **Calcium Signaling and β 2-Adrenergic Receptors Regulate 1-Nitropyrene Induced**
3 **CXCL8 Responses in BEAS-2B Cells**

4
5
6 Author list:

7 **Abdullah Mayati^{1,2*}, Eric Le Ferrec^{1,2*}, Jørn A. Holme³, Olivier Fardel^{1,2}, Dominique**
8 **Lagadic-Gossmann^{1,2}, Johan Øvrevik³**

9
10 ***Contributed equally to the publication**

11
12 Affiliation:

13 **¹UMR Inserm 1085, Institut de Recherche en Santé, Environnement, Travail, Rennes,**
14 **France**

15 **²Université de Rennes 1, SFR BIOSIT, Faculté des Sciences Pharmaceutiques et**
16 **Biologiques, Rennes, France.**

17 **³Department of Air Pollution and Noise, Division of Environmental Medicine,**
18 **Norwegian Institute of Public Health, Oslo, Norway**

19
20
21 Corresponding Author:

22 **Johan Øvrevik**

23 **Department of Air Pollution and Noise, Division of Environmental Medicine, Norwegian**
24 **Institute of Public Health, P.O. Box 4404 Nydalen, N-0403 Oslo, Norway**

25
26 **Phone: +47 21 07 64 08/Fax: +47 21 07 66 86**

27 **E-mail: johan.ovrevik@fhi.no**

Abstract

Nitro-polycyclic aromatic hydrocarbons (nitro-PAHs) are widespread environmental pollutants, generated from reactions between PAHs and nitrogen oxides during combustion processes. In the present study we have investigated the mechanisms of CXCL8 (IL-8) responses induced by 1-nitropyrene (1-NP) in human bronchial epithelial BEAS-2B cells, with focus on the possible importance of Ca^{2+} -signaling and activation of β_2 -adrenergic receptors ($\beta_2\text{AR}$). Ca^{2+} -chelator treatment obliterated 1-NP-induced CXCL8 (IL-8) responses. 1-NP at 10 μM (but not 1 μM) induced a rapid and sustained increase in intracellular Ca^{2+} -levels ($[\text{Ca}^{2+}]_i$). The early but not the later, sustained phase of 1-NP-induced $[\text{Ca}^{2+}]_i$ was suppressed by beta-blocker treatment (carazolol). Moreover, inhibition of $\beta_2\text{AR}$ by blocking-antibody, beta-blocker treatment (ICI 118551) or siRNA transfection attenuated CXCL8 responses induced by 1-NP. The results confirm that PAHs may induce Ca^{2+} -signaling also in BEAS-2B cells, at least partly through activation of $\beta_2\text{AR}$, and suggest that both $\beta_2\text{AR}$ - and Ca^{2+} -signaling may be involved in 1-NP-induced CXCL8 responses in bronchial epithelial cells.

Key words: Lung; Inflammation; Chemokines; Polycyclic Aromatic Hydrocarbons; Calcium; Adrenergic Receptors.

1

2 **Introduction**

3 Nitro-polycyclic aromatic hydrocarbons (nitro-PAHs) are ubiquitous air pollutants associated
4 with combustion particles, in particular diesel exhaust particles. Nitro-PAHs have long been
5 considered among the main contributors to the mutagenic effects of DEP (Hayakawa et al.,
6 1997; Scheepers et al., 1995). Nitro-PAHs may also exhibit considerable pro-inflammatory
7 potential, by inducing cytokine and chemokine responses in epithelial lung cells (Øvrevik et
8 al., 2010; Øvrevik et al., 2009; Park and Park, 2009; Pei et al., 2002).

9

10 Previous studies have shown that 1-nitropyrene (1-NP) and 3-nitrofluoranthene (3-NF) are
11 particularly potent inducers of the neutrophil attracting chemokine CXCL8 (IL-8), compared
12 to their amine counterparts (1-AP and 3-AF), un-substituted pyrene or benzo[a]pyrene
13 (B[a]P) (Øvrevik et al., 2010; Øvrevik et al., 2009; Øvrevik et al., 2013). This marked effect
14 of 1-NP and 3-NF on CXCL8 induction seems to be independent of aryl-hydrocarbon
15 receptor (AhR) activation, although it appears to involve formation of reactive oxygen species
16 (ROS) and/ or reactive electrophilic metabolites at least partly formed via CYP-mediated
17 metabolism (Øvrevik et al., 2010; Øvrevik et al., 2013).

18

19 Of notice, it has recently been shown that certain PAHs may induce calcium signaling in
20 endothelial cells through an aryl-hydrocarbon receptor (AhR)-independent mechanism
21 (Mayati et al., 2011). The effects were reported to be due to direct PAH-mediated activation
22 of β_2 -adrenergic receptors (β_2 AR), leading to a G protein/adenylyl cyclase/cyclic-AMP-
23 mediated calcium release from the endoplasmic reticulum (Mayati et al., 2012). Pyrene
24 seemed to be a particular potent inducer of this pathway and caused considerably stronger
25 calcium response in endothelial cells compared to other PAHs including B[a]P, chrysene and

benzo[*e*]pyrene (Mayati et al., 2011). Besides, we previously showed that the cell-permeable calcium chelator BAPTA-AM attenuated CXCL8 responses in BEAS-2B cells by a panel of environmental pollutants, including 1-NP (Øvrevik et al., 2011). Thus, in the present study we have investigated the 1-NP-induced effects on intracellular calcium levels in relation to CXCL8 responses in BEAS-2B cells, in order to test the hypothesis that β 2AR-mediated calcium-signaling may be a central mechanism for the pro-inflammatory effects of nitro-PAHs. Our results suggest that 1-NP is able to induce β 2AR-mediated calcium responses in BEAS-2B cells, and that both β 2AR- and Ca^{2+} -signaling are involved in 1-NP-induced CXCL8 responses. However, 1-NP also appeared to affect intracellular calcium levels through additional mechanisms, and β 2AR-mediated Ca^{2+} -signaling alone may not be sufficient for CXCL8-induction.

Materials and methods

Reagents

B[a]P, 1-NP, dimethyl sulphoxide (DMSO), ICI 118551 and carazolol were purchased from Sigma-Aldrich (St. Louis, MO, USA). LHC-9 cell culture medium and fura-2 acetoxymethyl ester (Fura-2-AM) were from Invitrogen (Carlsbad, CA, USA). Cytokine ELISA assay for CXCL8 (Human IL-8 Cytoset) was purchased from Biosource International (Camarillo, CA, USA). Antibodies against β 2AR (sc-569) were from Santa Cruz Biotechnology (Santa Cruz, CA, USA), whereas antibodies against β -actin were from Sigma-Aldrich. Short interfering RNA (siRNA) against β 2AR (sc-35147) with corresponding non-targeting control siRNA (sc-37007) were from Santa Cruz Biotechnology (CA, USA). All other chemicals used were purchased from commercial sources at the highest purity available.

1 *Cell cultures and exposures*

2 The BEAS-2B cell line, a SV40 hybrid (Ad12SV40) transformed human bronchial epithelial
3 cell line, was from European Collection of Cell Cultures (ECACC, Salisbury, UK). Cells
4 were maintained in LHC-9 medium in collagen-coated (PureCol™, Inamed Biomaterials,
5 Fremont, CA, USA) flasks in a humidified atmosphere at 37°C with 5% CO₂, and passaged
6 twice per week. Prior to exposure, cells were plated in 12-well culture dishes, grown to near
7 confluence in serum-free LHC-9 medium and exposed to PAHs and inhibitors as described
8 elsewhere. Controls for 1-NP -exposed cells were treated with vehicle (DMSO) only. DMSO
9 concentrations in all samples were below 0.5%.

11 *Chemokine release*

12 CXCL8 protein levels in cell-supernatants were determined by ELISA (Biosource
13 International, Camarillo, CA, USA), as described elsewhere (Øvrevik et al., 2010).
14 Absorbance was measured using a plate reader (TECAN Sunrise, Phoenix Research Products,
15 Hayward, CA, USA) complete with software (Magellan V 1.10).

17 *Gene silencing by siRNA*

18 Cells were reverse-transfected with siRNAs against β 2AR or non-targeting siRNAs, using
19 HiPerFect transfection reagent as recommended by the manufacturer (Qiagen, Germany: Fast-
20 Forward protocol for adherent cells). SiRNAs and HiPerFect were mixed by vortexing in
21 LHC-9 medium, incubated at room temperature (5-10 min) to form transfection complexes,
22 and added drop-wise to the cell cultures (100 μ l/well) immediately after seeding (at a final
23 siRNA concentration of 10 nM and 2.75 μ l of HiPerFect in a total of 1 ml growth medium).
24 Gene silencing was monitored by measuring β 2AR protein levels by Western blotting.

Calcium measurements

Variations in intracellular Ca^{2+} concentrations ($[\text{Ca}^{2+}]_i$) were analyzed by microspectrofluorimetry using the acetoxymethyl ester form of the Ca^{2+} -sensitive probe Fura-2, as previously reported (Le et al., 2002; N'Diaye et al., 2006). Cells were then submitted alternatively to 340 and 380 nm excitation wavelengths and the fluorescence from the trapped dye was measured at 510 nm. The F340/F380 ratio, i.e., ratio of fluorescence intensities after excitation at 340 nm and 380 nm, respectively, was used to estimate $[\text{Ca}^{2+}]_i$. To avoid potential problems with autofluorescence, PAH-concentrations were restricted to 10 μM in the $[\text{Ca}^{2+}]_i$ -assay.

Statistical analysis

Statistical significance was evaluated by GraphPad Prism software (GraphPad Software Inc., San Diego, CA, USA), using analysis of variance (ANOVA) with Bonferroni post-test.

Results

Role of calcium signaling in 1-NP-induced CXCL8 responses in BEAS-2B cells

We have previously shown that 1-NP induces a concentration-dependent increase in CXCL8 release from BEAS-2B cells from 10 μM (Øvrevik et al., 2010; Øvrevik et al., 2013). In line with this, we observed that 1-NP induced a strong increase in CXCL8 gene expression and protein release in BEAS-2B cells (Fig 1 A and B). To assess the role of Ca^{2+} -signaling in 1-NP-induced CXCL8 responses, BEAS-2B cells were pre-treated with the cell permeable Ca^{2+} -chelator BAPTA-AM for 30 min prior to 1-NP exposure for 18 h. BAPTA-AM completely blocked the 1-NP-induced CXCL8-response in BEAS-2B cells without affecting basal

CXCL8 levels (Fig. 1B). Thus, 1-NP-induced CXCL8 responses seemed to depend on intracellular Ca^{2+} -levels in BEAS-2B cells.

We next assessed whether 1-NP could affect intracellular Ca^{2+} -concentrations ($[\text{Ca}^{2+}]_i$) in BEAS-2B cells. Cells were loaded with the Ca^{2+} -sensitive fluorescent probe Fura-2 prior to 1-NP exposure, and changes in $[\text{Ca}^{2+}]_i$ were assessed by microspectrofluorimetry. As seen from Fig 2A, 10 μM 1-NP induced a time-dependent increase in $[\text{Ca}^{2+}]_i$ in BEAS-2B cells. The kinetics of 1-NP-induced $[\text{Ca}^{2+}]_i$ in BEAS-2B cells resembled that reported for B[a]P in endothelial HMEC cells (Mayati et al., 2011; Mayati et al., 2012) with a small, transient peak around 5 min after exposure, followed by a gradual increase that reached maximum after about half an hour exposure. Moreover, it has recently been reported that PAHs may induce increases in $[\text{Ca}^{2+}]_i$ in HMEC cells through direct activation of $\beta 2\text{AR}$ (Mayati et al., 2012). Thus, to assess whether similar mechanisms could be involved in the present PAH-induced Ca^{2+} response, BEAS-2B cells were pre-incubated with the β -blocker carazolol for 30 min prior to exposure with 10 μM 1-NP. Carazolol almost completely blocked the early phases of 1-NP induced increases in $[\text{Ca}^{2+}]_i$ (Fig. 2B). However, 1-NP induced a sustained increase in $[\text{Ca}^{2+}]_i$ lasting at least up to 6 h (Fig. S1A, online supplementary materials). The later phases of 1-NP-induced (from 45 min to 6 h) seemed unaffected by Carazolol treatment (Fig. 2B).

Of notice, previous studies have shown that PAHs such as B[a]P and un-substituted pyrene may stimulate $\beta 2\text{AR}$ -mediated $[\text{Ca}^{2+}]_i$ already at 1 μM concentration (Mayati et al., 2011; Mayati et al., 2012). In line with this we observed that 1 μM B[a]P was able to induce increased $[\text{Ca}^{2+}]_i$ in BEAS-2B cells. However, 1-NP failed to affect Ca^{2+} signaling at this low concentration. Moreover, at 10 μM B[a]P appeared to induce considerably higher effects on $[\text{Ca}^{2+}]_i$ compared to 1-NP (Fig S1B and C, online supplementary materials). Thus, although 1-

NP was able to stimulate $\beta 2AR$ -induced $[Ca^{2+}]_i$, it appeared to be a less potent activator compared to other PAHs.

Role of $\beta 2AR$ in 1-NP-induced CXCL8 responses in BEAS-2B cells

Next we wanted to assess whether $\beta 2AR$ could be involved in nitro-PAH-induced CXCL8 responses in BEAS-2B cells. Therefore we pre-incubated the cells with a $\beta 2AR$ -blocking antibody (Mayati et al., 2012) for 30 min prior to exposure with 1-NP. The $\beta 2AR$ -blocking antibody suppressed 1-NP-induced CXCL8 responses by approximately 50% at both tested concentrations (Fig 3A). To further examine the role of $\beta 2AR$ in 1-NP-induced CXCL8 responses, we assess the effects of the selective $\beta 2AR$ -antagonist ICI 118551 and of silencing $\beta 2AR$ by siRNA (si $\beta 2AR$). Both ICI treatment and transfection with si $\beta 2AR$ resulted in a partial, but statistically significant suppression of 1-NP-induced CXCL8 (Fig 3B and C), comparable to the effect obtained with the $\beta 2AR$ -blocking antibody. Thus $\beta 2AR$ -signaling seemed to be involved in 1-NP-induced chemokine responses.

Discussion

In the present study we have assessed the mechanisms of the 1-NP induced CXCL8 response with emphasis on the importance of Ca^{2+} -signaling and the $\beta 2AR$ -receptor. Previously, Ca^{2+} -chelation was found to suppress CXCL8 in BEAS-2B cells by a variety of different air pollution constituents including 1-NP (Øvrevik et al., 2011). In coherence with this, the present results confirmed that the cell-permeable Ca^{2+} -chelator BAPTA-AM almost completely block 1-NP-induced CXCL8. Of interest, PAHs may increase $[Ca^{2+}]_i$ in endothelial cells through AhR-independent mechanisms (Mayati et al., 2011). Recent findings suggest that this effect may be regulated through PAH-mediated binding and activation of the

1 β 2AR (Mayati et al., 2012). In support of this, we observed that 1-NP (10 μ M) exposure
2 induced an increase in $[Ca^{2+}]_i$ in BEAS-2B cells that could be suppressed by beta-blocker
3 treatment. Thus, PAH-induced Ca^{2+} -signaling through β 2AR-activation may extend to several
4 cell types. However, only the early phase of the Ca^{2+} -response seemed to depend on signaling
5 through β -adrenergic receptors, as beta-blocker treatment had no effect on the 1-NP induced
6 increase in $[Ca^{2+}]_i$ after 40 min exposure. Thus it is conceivable that 1-NP may induce Ca^{2+} -
7 signaling through multiple mechanisms.

8
9 Interference with β 2AR-signalling by use of blocking antibodies, a pharmacological beta-
10 blocker, or by silencing β 2AR-expression by siRNA-transfection, led to a suppression of 1-
11 NP-induced CXCL8 responses. Thus 1-NP-induced CXCL8 release in BEAS-2B cells
12 seemed at least partly dependent on β 2AR-activation. In line with this, the β 2AR agonists
13 salbutamol and salmeterol have been found to enhance CXCL8 and interleukin-6 (IL-6)
14 responses by IL-1 β or virus infections in BEAS-2B cells and primary human bronchial
15 epithelial cells (Edwards et al., 2007; Holden et al., 2010). However, salbutamol and
16 salmeterol had no effect on CXCL8 or IL-6 responses alone, suggesting that β 2AR-signalling
17 alone may be insufficient for activation of cytokine/chemokine responses. Indeed, B[a]P
18 appeared to be a more potent inducer of β 2AR-dependent $[Ca^{2+}]_i$ -responses than 1-NP, but is
19 nevertheless unable to induce CXCL8 or other chemokines in BEAS-2B cells (Øvrevik et al.,
20 2010; Øvrevik et al., 2013). Thus, additional 1-NP-induced mechanisms are likely required
21 for the response. This notion is supported by previous findings suggesting that 1-NP-induced
22 CXCL8 responses also involve metabolic activation by CYP-enzymes and possibly ROS
23 (Øvrevik et al., 2013).

24

While β 2AR-interference only partly attenuated 1-NP-induced CXCL8 release, Ca^{2+} -chelation by BAPTA-AM completely abrogated the CXCL8 response. It should therefore be considered that the role of β 2AR in 1-NP-induced CXCL8-release could be linked to other signaling mechanisms such as cAMP or β -arrestin. Indeed, salbutamol-induced exacerbation of virus-induced IL-6 responses in BEAS-2B cells seemed to be mediated through a cAMP-dependent mechanism (Edwards et al., 2007). If so, the effects of BAPTA-AM on 1-NP-induced CXCL8 may be also related to interference with the later, sustained, β 2AR-independent $[\text{Ca}^{2+}]_i$ -response.

The strong CXCL8-induction by 1-NP seems predominately to occur at high concentrations ($\geq 10 \mu\text{M}$) in BEAS-2B cells (Øvrevik et al., 2013). However, we recently observed that low concentrations of 1-NP ($1 \mu\text{M}$; giving no cytokine release alone) potentiated CXCL8-responses induced by priming the cells with a Toll-like receptor 3-agonist. Similar effects were observed with low concentrations of 1-AP and un-substituted pyrene (Øvrevik et al., 2013). Since 1-NP may exacerbate CXCL8 responses at concentrations that are insufficient to induce β 2AR-dependent $[\text{Ca}^{2+}]_i$, other cellular targets seem to exist that are more sensitive towards 1-NP than the β 2AR. Such low-concentration targets may be more important scenarios of real-life exposure to pyrene and pyrene-derivatives. Other PAHs may have considerably higher affinity for β 2AR. As reported in the present study and elsewhere (Mayati et al., 2012), B[a]P may induce β 2AR-dependent $[\text{Ca}^{2+}]_i$ -responses already at $1 \mu\text{M}$. In fact, the affinity of B[a]P ($K_d = 10 \text{ nM}$) appears to be among the highest reported for β 2AR ligands (Mayati et al., 2012). However, it should also be noted that 1-NP concentrations in diesel exhaust particles may be up to 10-fold higher than the concentration of B[a]P (Totlandsdal et al., 2012; Totlandsdal et al., 2014).

In extension of previous observations from endothelial cells (Mayati et al., 2011; Mayati et al., 2012), the present results suggest that PAHs may induce increased $[Ca^{2+}]_i$ through $\beta 2AR$ -activation also in epithelial lung cells. Moreover, $\beta 2AR$ -signalling and $[Ca^{2+}]_i$ appeared to be involved in the regulation of 1-NP-induced CXCL8 in the BEAS-2B cells. Although other yet unidentified cellular targets may be more sensitive to 1-NP exposure, $\beta 2AR$ appears to be a highly sensitive for other PAHs (Mayati et al., 2012). PAHs have also been reported to impair $\beta 2AR$ -function and interfere with asthma treatment (Factor et al., 2011). Thus, further studies are warranted to clarify the role of $\beta 2AR$ in PAH-induced responses and its possible implications in lung toxicity.

Acknowledgements

The work was supported by the Research Council of Norway, through the Environmental Exposures and Health Outcomes-program (grants no. 185620 and 228143).

Reference List

- Edwards, M.R., Haas, J., Panettieri, R.A., Jr., Johnson, M., Johnston, S.L., 2007. Corticosteroids and beta2 agonists differentially regulate rhinovirus-induced interleukin-6 via distinct Cis-acting elements. *J Biol Chem* 282, 15366-15375.
- Factor, P., Akhmedov, A.T., McDonald, J.D., Qu, A., Wu, J., Jiang, H., Dasgupta, T., Panettieri, R.A., Jr., Perera, F., Miller, R.L., 2011. Polycyclic aromatic hydrocarbons impair function of beta2-adrenergic receptors in airway epithelial and smooth muscle cells. *Am J Respir. Cell Mol Biol* 45, 1045-1049.
- Hayakawa, K., Nakamura, A., Terai, N., Kizu, R., Ando, K., 1997. Nitroarene concentrations and direct-acting mutagenicity of diesel exhaust particulates fractionated by silica-gel column chromatography. *Chem. Pharm Bull. (Tokyo)* 45, 1820-1822.
- Holden, N.S., Rider, C.F., Bell, M.J., Velayudhan, J., King, E.M., Kaur, M., Salmon, M., Giembycz, M.A., Newton, R., 2010. Enhancement of inflammatory mediator release by beta(2)-adrenoceptor agonists in airway epithelial cells is reversed by glucocorticoid action. *Br J Pharmacol* 160, 410-420.

- 1 Le, F.E., Lagadic-Gossmann, D., Rauch, C., Bardiau, C., Maheo, K., Massiere, F., Le, V.M.,
2 Guillouzo, A., Morel, F., 2002. Transcriptional induction of CYP1A1 by oltipraz in
3 human Caco-2 cells is aryl hydrocarbon receptor- and calcium-dependent. *J Biol*
4 *Chem* 277, 24780-24787.
- 5 Mayati, A., Le Ferrec, E., Lagadic-Gossmann, D., Fardel, O., 2011. Aryl hydrocarbon
6 receptor-independent up-regulation of intracellular calcium concentration by
7 environmental polycyclic aromatic hydrocarbons in human endothelial HMEC-1 cells.
8 *Environ.Toxicol.*
- 9 Mayati, A., Levoin, N., Paris, H., N'Diaye, M., Courtois, A., Uriac, P., Lagadic-Gossmann,
10 D., Fardel, O., Le, F.E., 2012. Induction of intracellular calcium concentration by
11 environmental benzo(a)pyrene involves a beta2-adrenergic receptor/adenylyl
12 cyclase/Epac-1/inositol 1,4,5-trisphosphate pathway in endothelial cells. *J Biol Chem*
13 287, 4041-4052.
- 14 N'Diaye, M., Le, F.E., Lagadic-Gossmann, D., Corre, S., Gilot, D., Lecureur, V., Monteiro,
15 P., Rauch, C., Galibert, M.D., Fardel, O., 2006. Aryl hydrocarbon receptor- and
16 calcium-dependent induction of the chemokine CCL1 by the environmental
17 contaminant benzo[a]pyrene. *J Biol Chem* 281, 19906-19915.
- 18 Øvrevik, J., Arlt, V.M., Oya, E., Nagy, E., Møllerup, S., Phillips, D.H., Lag, M., Holme, J.A.,
19 2010. Differential effects of nitro-PAHs and amino-PAHs on cytokine and chemokine
20 responses in human bronchial epithelial BEAS-2B cells. *Toxicol.Appl.Pharmacol.*
21 242, 270-280.
- 22 Øvrevik, J., Lag, M., Holme, J.A., Schwarze, P.E., Refsnes, M., 2009. Cytokine and
23 chemokine expression patterns in lung epithelial cells exposed to components
24 characteristic of particulate air pollution. *Toxicology* 259, 46-53.
- 25 Øvrevik, J., Refsnes, M., Holme, J.A., Schwarze, P.E., Lag, M., 2013. Mechanisms of
26 chemokine responses by polycyclic aromatic hydrocarbons in bronchial epithelial
27 cells: sensitization through toll-like receptor-3 priming. *Toxicol Lett.* 219, 125-132.
- 28 Øvrevik, J., Refsnes, M., Totlandsdal, A.I., Holme, J.A., Schwarze, P.E., Lag, M., 2011.
29 TACE/TGF- α /EGFR regulates CXCL8 in bronchial epithelial cells exposed to
30 particulate matter components. *Eur Respir.J* 38, 1189-1199.
- 31 Park, E.J., Park, K., 2009. Induction of pro-inflammatory signals by 1-nitropyrene in cultured
32 BEAS-2B cells. *Toxicol.Lett.* 184, 126-133.
- 33 Pei, X.H., Nakanishi, Y., Inoue, H., Takayama, K., Bai, F., Hara, N., 2002. Polycyclic
34 aromatic hydrocarbons induce IL-8 expression through nuclear factor kappaB
35 activation in A549 cell line. *Cytokine* 19, 236-241.
- 36 Scheepers, P.T., Martens, M.H., Velders, D.D., Fijneman, P., van Kerkhoven, M., Noordhoek,
37 J., Bos, R.P., 1995. 1-Nitropyrene as a marker for the mutagenicity of diesel exhaust-
38 derived particulate matter in workplace atmospheres. *Environ.Mol.Mutagen.* 25, 134-
39 147.

- 1 Totlandsdal, A.I., Herseth, J.I., Bolling, A.K., Kubatova, A., Braun, A., Cochran, R.E.,
2 Refsnes, M., Øvrevik, J., Lag, M., 2012. Differential effects of the particle core and
3 organic extract of diesel exhaust particles. *Toxicol Lett.* 208, 262-268.
- 4 Totlandsdal, A.I., Øvrevik, J., Cochran, R.E., Herseth, J.I., Bolling, A.K., Lag, M., Schwarze,
5 P., Lilleaas, E., Holme, J.A., Kubatova, A., 2014. The occurrence of polycyclic
6 aromatic hydrocarbons and their derivatives and the proinflammatory potential of
7 fractionated extracts of diesel exhaust and wood smoke particles. *J Environ Sci Health*
8 *A Tox.Hazard.Subst.Environ Eng* 49, 383-396.
9
10
11

Figure legends


FIGURE 1. CXCL8 gene expression and protein release in 1-NP-exposed BEAS-2B cells, and role of Ca^{2+} -chelation by BAPTA-AM. Cells were exposed to 20 μM 1-NP or vehicle (DMSO) alone. CXCL8 gene expression was measured after 6 h by real-time PCR (A). Cells were pre-incubated with 10 μM of the cell-permeable Ca^{2+} -chelator BAPTA-AM for 30 min prior to exposure to 20 μM 1-NP or vehicle (DMSO) alone for 18 h (B). The figure depicts mean \pm SEM of three independent experiments. *Significantly different from unexposed controls ($P < 0.05$); †Significant down-regulation by chelator-treatment ($P < 0.05$).

FIGURE 2. Intracellular Ca^{2+} -levels and role of $\beta 2\text{AR}$ -signaling in 1-NP exposed BEAS-2B cells. Cells were pre-incubated with 10 μM of the β -blocker carazolol (Cara) prior to incubation with 10 μM 1-NP or vehicle (DMSO) alone (A and B). Intracellular Ca^{2+} -concentrations ($[\text{Ca}^{2+}]_i$) were measured by incubation with the Ca^{2+} -sensitive probe Fura-2 and the ratio of fluorescence intensities after excitation at 340 nm and 380 nm, respectively, was used to estimate $[\text{Ca}^{2+}]_i$. Figure A depicts the mean of normalized $[\text{Ca}^{2+}]_i$ from two independent continuous recordings, while figure B depicts the mean \pm SEM of three independent experiments after 6 h (B) exposure.


FIGURE 3. Role of $\beta 2\text{AR}$ -signaling in CXCL8-responses in 1-NP-exposed BEAS-2B cells. Cells were pre-incubated for 30 min with a $\beta 2\text{AR}$ -blocking antibody (A) or the β -blocker ICI 118551 (B), or transfected with siRNA against $\beta 2\text{AR}$ (si $\beta 2\text{AR}$) or non-targeting control siRNA (siNT) (C), prior to exposure with 20 μM 1-NP or vehicle (DMSO) alone for 18 h. CXCL8 release were measured by ELISA. Efficiency of $\beta 2\text{AR}$ knock-down by siRNA

1 was assessed by Western blotting (C). The figures depict mean \pm SEM of three or more
2 independent experiments. *Significantly different from unexposed controls ($P < 0.05$);
3 †Significant down-regulation by inhibitor/antibody/siRNA treatment ($P < 0.05$).
4


ACCEPTED MANUSCRIPT




A



B





1

2 **Highlights**

- 3 • We examined mechanisms of 1-NP induced CXCL8 responses in BEAS-2B cells
- 4 • Treatment with a Ca^{2+} -chelator abrogated the 1-NP-induced CXCL8 response
- 5 • 1-NP induced a rapid increase in intracellular Ca^{2+} -levels
- 6 • Beta-blocker treatment suppressed the 1-NP-induced Ca^{2+} -responses
- 7 • Inhibition of $\beta 2\text{AR}$ suppressed 1-NP-induced CXCL8 responses

8

ACCEPTED MANUSCRIPT